

# Collapse of proteostasis represents an early molecular event in *Caenorhabditis elegans* aging

Anat Ben-Zvi, Elizabeth A. Miller, and Richard I. Morimoto<sup>1</sup>

Department of Biochemistry, Molecular Biology and Cell Biology, Rice Institute for Biomedical Research, Northwestern University, Evanston, IL 60208

Edited by Susan L. Lindquist, Whitehead Institute for Biomedical Research, Cambridge, MA, and approved July 22, 2009 (received for review March 15, 2009)

**Protein damage contributes prominently to cellular aging. To address whether this occurs at a specific period during aging or accumulates gradually, we monitored the biochemical, cellular, and physiological properties of folding sensors expressed in different tissues of *C. elegans*. We observed the age-dependent misfolding and loss of function of diverse proteins harboring temperature-sensitive missense mutations in all somatic tissues at the permissive condition. This widespread failure in proteostasis occurs rapidly at an early stage of adulthood, and coincides with a severely reduced activation of the cytoprotective heat shock response and the unfolded protein response. Enhancing stress responsive factors HSF-1 or DAF-16 suppresses misfolding of these metastable folding sensors and restores the ability of the cell to maintain a functional proteome. This suggests that a compromise in the regulation of proteostatic stress responses occurs early in adulthood and tips the balance between the load of damaged proteins and the proteostasis machinery. We propose that the collapse of proteostasis represents an early molecular event of aging that amplifies protein damage in age-associated diseases of protein conformation.**

daf-16 | folding sensors | hsf-1 | protein misfolding | stress response

The stability of the proteome is challenged by conditions that cause proteotoxic stress including errors during protein synthesis, oxidant-induced covalent modifications, inherited polymorphisms, and misfolding (1–3). Consequently, all cells have highly conserved stress-inducible pathways that detect, prevent, and resolve such damage (4, 5). Accumulation of misfolded proteins titrate the negative regulation of chaperones from HSF-1 (4, 6, 7). Likewise, an E3 ubiquitin ligase was shown to regulate the levels of DAF-16, also indicating a link between stress response activation to the balance between proteostasis capacity and the load of damaged proteins (8). The activation of cellular stress responses limits the accumulation of damaged proteins by suppression of misfolding and enhancing chaperone-mediated folding, promoting the enzymatic removal of covalent modifications, and stimulating clearance by ubiquitin-mediated, proteosomal, and autophagic processes (1, 4, 5, 9, 10). The dynamics of these processes during aging, however, are poorly understood.

Aging can be characterized by the loss of cellular function and increased vulnerability to environmental and physiological stress that results in enhanced susceptibility to disease. Many age-dependent changes have been detected at mid-to-late lifespan of *C. elegans*, including the accumulation of damaged macromolecules, cell and organellar degeneration, and physiological decline (11–17). Likewise, aggregation and toxicity of disease-associated aggregation-prone proteins, such as huntingtin and A $\beta$  peptide, is enhanced during aging in *C. elegans* and other model systems (18–21).

The regulation of aging and the maintenance of proteostasis has been recently established in *C. elegans*. Downregulation of the insulin-like signaling (ILS) pathway enhances lifespan and stress resistance, and suppresses aggregation of polyglutamine expansion proteins and A $\beta$  toxicity (20–22). This requires the activities of the stress responsive factors HSF-1 and DAF-16, and

the elevated expression of molecular chaperones (20, 21, 23–26). These observations have led to the hypothesis that there is an age-dependent failure in protein-folding quality control. Whether damaged proteins are functionally sequestered and neutral to cellular processes, toxic, or cleared more rapidly at different stages of development and adulthood is likely to have profound implications for the functionality of biological systems when challenged by aging and disease.

To examine the stability of the proteome during aging, we monitored the activities of folding sensors that harbor missense mutations with temperature-sensitive (ts) phenotypes. We selected diverse, well-characterized ts mutations expressed in different tissues that are fully functional under physiological permissive (15 °C) conditions and exhibit a complete loss of function at the restrictive (25 °C) conditions. Here we show that each folding sensor misfolds at an early stage of adulthood during a period that coincides with a sharp decline in the ability to activate cellular stress responses. This collapse of proteostasis can be restored by the stress responsive factors HSF-1 and DAF-16, thus prolonging the health of the organism. Dysregulation of proteostasis at the transition to adulthood offers an explanation for the age-dependent onset observed for many heritable diseases.

## Results

The effects of aging on proteostasis in *C. elegans* muscle cells were examined using ts mutations in the thick filament proteins paramyosin (UNC-15) and myosin heavy chain B (UNC-54). Animals expressing paramyosin(ts) [*unc-15(e1402)*] or myosin(ts) [*unc-54(e1301)*] at the restrictive condition exhibit severe movement defects (uncoordination) and have disrupted thick filaments that result from misfolding of paramyosin(ts) or myosin(ts) (2, 27, 28). At the permissive temperature, as expected, paramyosin(ts)- or myosin(ts)-expressing animals show wild-type (WT) phenotypes through day  $\approx$ 6 of adulthood (Fig. 1 *A* and *G*). However, at day  $\approx$ 7 of adulthood and thereafter, animals expressing either ts-mutant protein exhibited significant uncoordination, whereas the motility of WT animals was unaffected until day 10 of adulthood, when aging-associated decline in motility was observed (Fig. 1 *A* and *G*) (11).

To determine whether the decline in motility, observed for animals expressing muscle cell ts-mutant proteins, is due to disruption of myofilaments, we used cellular and biochemical assays to monitor the subcellular localization of paramyosin(ts) and myosin(ts). WT animals expressing myosin heavy chain A tagged with green fluorescence protein (GFP) (MYO-3::GFP) provided a complementary assay to monitor the state of myofilament structure. Animals expressing either ts-mutant protein


Author contributions: A.B.-Z. and R.I.M. designed research; A.B.-Z. and E.A.M. performed research; A.B.-Z. analyzed data; and A.B.-Z. and R.I.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: r-morimoto@northwestern.edu.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0902882106/DCSupplemental](http://www.pnas.org/cgi/content/full/0902882106/DCSupplemental).



**Fig. 1.** The folding and assembly of metastable proteins in muscle cells are perturbed in early adulthood. (A) Percentage of slow-moving, age-synchronized animals expressing WT (squares) or ts (triangles) paramyosin (*unc-15*) at 15 °C. Data are mean ± SD, >95 animals per data point. (B) Percent of cells with disrupted myofilaments, quantified from confocal images of MYO-3::GFP fluorescence of animals expressing WT (squares) or ts (triangles) paramyosin at 15 °C. Data are mean ± SD, >210 cells per data point. (C) Differential sensitivity of paramyosin to chymotrypsin proteolysis. Total protein extracts of age-synchronized animals were treated for indicated times, and gels were incubated with paramyosin (*Top*) or dynamin (as an internal control) (*Bottom*) antibodies. Proteolysis experiments were performed on 3 independent biological samples. Images are from a representative proteolysis experiment. (D) Quantification of proteolyzed paramyosin (fragment 3 + 4) (black) and dynamin (gray) after 1 h of treatment normalized to day 1. (E and F) Confocal images of body-wall muscles. Age-synchronized animals expressing WT or ts paramyosin grown at 15 °C (E) or 25 °C (F) were stained with anti-paramyosin and phalloidin. (Scale bar: 5 μm.) (G) Percentage of slow movement of animals expressing WT (squares) or ts (triangles) myosin (*unc-54*) as in (A). Data are mean ± SD, >110 animals per data point. (H) Percent of cells with disrupted myofilaments expressing WT (squares) or (triangles) ts myosin as in (B). Data are mean ± SD, >190 cells per data point. (I) Differential sensitivity of myosin(ts) protein to enterokinase digestion as in (C). Gels were exposed to myosin (*Top*) or dynamin (*Bottom*) antibodies. (J) Quantification of digested myosin (fragment 1 levels after 7 h digest) (black) or dynamin (gray) from (I) normalized to day 1. (K and L) Confocal images of animals expressing WT or ts myosin grown at 15 °C (K) or 25 °C (L) as in (E). Labels: wt, wild type (WT); ts, time-sensitive.





occurs early in adulthood. Relative to other biological markers of aging that have identified changes at the midpoint of lifespan of *C. elegans*, at days 8–14 of adulthood, the loss of proteostasis occurs at a much earlier stage in multiple cell types, including neurons and muscles (11–17). How does an early event in adulthood affect cellular pathology and loss of physiological function during aging? Our findings show that aging-associated sarcopenia, a general biomarker of aging, corresponds to misfolding of various WT muscle proteins. Given that muscle is a multiprotein complex that requires both specialized and general components of the proteostasis machinery for folding and assembly, it is possible that the decline in proteostasis capacity decreases muscle filaments stability (40). This interpretation is supported by the “disposable soma” theory of aging, which suggests that an early event of maintenance dysregulation can have profound downstream consequences on multiple cellular pathways and physiological functions later in life, as the accumulation of misfolded and damaged proteins is amplified during aging (41). This is further supported by the finding that the loss of proteostasis in early adulthood corresponds to the same period during adult aging when polyglutamine-expansion proteins exhibit age-dependent aggregation and toxicity in *C. elegans* (20). It is interesting to note that the lifespan for WT *C. elegans* under laboratory conditions is 2–3 weeks, whereas in soil conditions this was reduced by nearly an order of magnitude (42). This would suggest that the maintenance of proteostasis is essential throughout development and extends only through the reproductive phase of adulthood, when there is an evolutionary advantage to allocate resources to proteostasis maintenance (41).

The ability of HSF-1 and DAF-16 to restore proteostasis supports the central roles of these stress-responsive transcription factors in more general aspects of biology to prevent global instability of the proteome. Enhancing the activities of these master regulators during development and early in adulthood could affect the levels of proteostasis machinery, preventing proteostasis collapse (43, 44). This is supported by their central role in delaying aging-associated cellular dysfunction and age-related diseases (20, 21, 23, 24). Moreover, because both HSF-1 and DAF-16 are regulated by the NAD-dependent sirtuin, SIRT1, and the ILS-pathway, it is reasonable to suggest that stress biology and proteostasis are intimately linked to each other and to metabolic control and lifespan (20, 23, 24, 45–47). In *C. elegans*, ILS-dependent life extension and the activation of the HS response are regulated by active neuronal signaling and cell nonautonomous control (48, 49). We therefore suggest that the transition between development and adulthood is critical at the cellular and organismal levels to the regulation of proteostasis. Regulatory override of the switch by enhancing HSF-1 and DAF-16 could prolong the “youthfulness” of the proteostatic machineries and ensure that the regulation of proteostasis remains robust, thus reducing the rate of aging, and extending lifespan.

Our experiments, by their design, provide a semiquantitative analysis of specific phenotypes associated with each folding sensor. It seems reasonable to extend our observations to the effects of aging on the stability of other coexpressed metastable proteins, many of which are tissue specific and may lose function as the proteostasis network becomes compromised during aging. By extrapolation to the entire proteome, missense mutations are not rare, as 60,000 single nucleotide substitutions are estimated to occur in human coding exons (an average of 2 sites per gene) (50), and 25%–30% of these are predicted to affect protein stability or folding (51). Moreover, missense mutations represent only one source of damage that could affect the stability of the proteome. For example, misincorporation during translation is estimated to result in one missense substitution in 18% of expressed proteins, with additional opportunities to express damaged proteins from errors in transcrip-

tion and splicing, posttranslational modifications, and targeting (3). The spectrum of affected proteins would therefore be predicted to contribute significantly to the high degree of variability in onset and progression of aging-associated decline (11). The exposure of missense mutations during aging, which are normally masked by the proteostasis network, offers an explanation for the increased risk observed for many heritable age-dependent diseases, such as familial ALS, Parkinson’s disease, and Huntington’s disease (52–56).

## Materials and Methods

**Nematode Strains and Maintenance.** For a list of strains used in this work and name abbreviations see Tables S1 and S2. Nematodes were grown on NGM plates seeded with *Escherichia coli* OP50 strain at 15 °C. Animals were picked as L4 larvae and scored at the indicated times during adulthood. All animals were scored at the same chronological age early in adulthood. Animals were moved every 2–3 days during the reproductive period to avoid progeny contamination, and were discarded after scoring. Experiments were repeated at least 3 times. All assays were performed blind.

**Assays for Specific Temperature-Sensitive Phenotypes.** For the Osm phenotype, slow movement (Unc), paralysis, and defective ssGFP uptake assays, animals were scored as previously described (2, 30). For sensitivity to EtOH or levamisole, animals were picked into 0.4 M EtOH or 50  $\mu$ M levamisole solution respectively, equilibrated for 5 min, and scored for motility. For MYO-3::GFP and SNB-1::GFP mislocalization, synchronized adults expressing the transgene were fixed (2) and GFP fluorescence was monitored. Animals were imaged using a Zeiss ConfoCor 3/510 META confocal microscope through a 63  $\times$  1.0 numerical aperture objective with a 488 nm line for excitation.

**Immunostaining.** Immunofluorescence studies were performed as previously described (2). Animals were stained with rhodamine phalloidin (Molecular Probes) and with either anti-paramyosin [5-23] (57), anti-myosin heavy chain B [28.2] (57), anti-myosin heavy chain A [5-6] (57) or anti-dynamin-1 antibodies (29). Secondary antibodies were IgG-FITC goat anti-mouse (Sigma) or Rhodamine Red goat anti-rabbit (Invitrogen). Animals were imaged as described using 488- and 543-nm lines for excitation.

**Proteolysis.** Nematodes were lysed, and digests were analyzed as previously described (2). Total protein of 120–200  $\mu$ g was digested at RT with chymotrypsin (Sigma) or enterokinase (Roche) at enzyme-to-protein ratio 1:1,200 and 1:33, respectively. Aliquots were removed at the indicated times. Each digest was repeated at least 3 times using independent biological samples. Protein levels were determined using ImageJ software. To quantify the change in fragmentation, we identified differences in the fragmentation pattern of animals grown at 15 °C or 25 °C (fragments 3 and 4 for paramyosin(ts) [see Fig. 1C] and fragment 1 for myosin(ts) [see Fig. 1I]) and examined the levels of these fragments after a digest of 1 h for paramyosin or 7 h for myosin, relative to undigested proteins of animals collected at indicated times during adulthood. Values were normalized to day 1 of adulthood.

**RNA Interference (RNAi).** Between 15 and 30 L4 larvae were placed on *E. coli* strain HT115(DE3) transformed with indicated RNAi vectors (J. Ahinger, University of Cambridge, Cambridge, U.K.) as previously described (58).

**RNA Levels.** Total RNA was extracted using TRIzol reagent. For cDNA synthesis, mRNA was reverse transcribed using the iScript™ cDNA Synthesis Kit (Bio-Rad). Quantitative PCR was performed as previously described (49). See Table S3 for primer sequences.

**Stress Induction.** A total of 15–20 age-synchronized WT animals were used for each assay. For HS, animals were moved to new plates, sealed, and placed at 32 °C for 30 min. For induction of UPR, animals were shifted to 24-well plates and grown for 7 h in M9 buffer supplemented with bacteria (final OD<sub>600</sub> 0.1) with or without 3 mM DTT. Approximately 15 animals were collected for each condition and frozen immediately following the stress.

**Thermotolerance Assay.** Animals were heat shocked at 35 °C for 6 h, and animals’ survival was examined by touch-induced movement (49).

**Lifespan.** Assays were performed at 15 °C as previously described (24). As was previously reported for other cultivation temperatures, overexpression of HSF-1 or DAF-16 increased the lifespan of animals grown at 15 °C by 15%–40% (24, 59).

**ACKNOWLEDGMENTS.** We thank Dr. Ilya Ruvinsky and members of the Morimoto lab for critical discussions and reading of the manuscript. We thank the Caenorhabditis Genetics Center, which is funded by the National Institutes of Health National Center for Research Resources for some of the nematode strains. Antibodies 5-6 and 5-23 were obtained from the Developmental

Studies Hybridoma Bank developed under the auspices of the National Institute of Child Health and Human Development and maintained by the University of Iowa, Department of Biological Sciences. These studies were supported by grants from the National Institutes of Health (National Institute of General Medical Sciences and National Institute on Aging).

- Stadtman ER (1992) Protein oxidation and aging. *Science* 257(5074):1220–1224.
- Gidalevitz T, Ben-Zvi A, Ho KH, Brignull HR, Morimoto RI (2006) Progressive disruption of cellular protein folding in models of polyglutamine diseases. *Science* 311(5766):1471–1474.
- Drummond DA, Wilke CO (2008) Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. *Cell* 134(2):341–352.
- Morimoto RI (2008) Proteotoxic stress and inducible chaperone networks in neurodegenerative disease and aging. *Genes Dev* 22(11):1427–1438.
- Ron D, Walter P (2007) Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8(7):519–529.
- Shi Y, Mosser DD, Morimoto RI (1998) Molecular chaperones as HSF1-specific transcriptional repressors. *Genes Dev* 12(5):654–666.
- Zou J, Guo Y, Guettouche T, Smith DF, Voellmy R (1998) Repression of heat shock transcription factor HSF1 activation by HSP90 (HSP90 complex) that forms a stress-sensitive complex with HSF1. *Cell* 94(4):471–480.
- Li W, Gao B, Lee SM, Bennett K, Fang D (2007) RLE-1, an E3 ubiquitin ligase, regulates C. elegans aging by catalyzing DAF-16 polyubiquitination. *Dev Cell* 12(2):235–246.
- Medicherla B, Goldberg AL (2008) Heat shock and oxygen radicals stimulate ubiquitin-dependent degradation mainly of newly synthesized proteins. *J Cell Biol* 182(4):663–673.
- Mizushima N, Levine B, Cuervo AM, Klionsky DJ (2008) Autophagy fights disease through cellular self-digestion. *Nature* 451(7182):1069–1075.
- Herndon LA, et al. (2002) Stochastic and genetic factors influence tissue-specific decline in ageing C. elegans. *Nature* 419(6909):808–814.
- Huang C, Xiong C, Kornfeld K (2004) Measurements of age-related changes of physiological processes that predict lifespan of Caenorhabditis elegans. *Proc Natl Acad Sci USA* 101(21):8084–8089.
- Garigan D, et al. (2002) Genetic analysis of tissue aging in Caenorhabditis elegans: A role for heat-shock factor and bacterial proliferation. *Genetics* 161(3):1101–1112.
- Haitcock E, et al. (2005) Age-related changes of nuclear architecture in Caenorhabditis elegans. *Proc Natl Acad Sci USA* 102(46):16690–16695.
- Yasuda K, et al. (2006) Age-related changes of mitochondrial structure and function in Caenorhabditis elegans. *Mech Ageing Dev* 127(10):763–770.
- Ishii N, Goto S, Hartman PS (2002) Protein oxidation during aging of the nematode Caenorhabditis elegans. *Free Radic Biol Med* 33(8):1021–1025.
- Gerstbrein B, Stamatas G, Kollias N, Driscoll M (2005) In vivo spectrofluorimetry reveals endogenous biomarkers that report healthspan and dietary restriction in Caenorhabditis elegans. *Aging Cell* 4(3):127–137.
- Davies SW, et al. (1997) Formation of neuronal intranuclear inclusions underlies the neurological dysfunction in mice transgenic for the HD mutation. *Cell* 90(3):537–548.
- Warrick JM, et al. (1998) Expanded polyglutamine protein forms nuclear inclusions and causes neural degeneration in Drosophila. *Cell* 93(6):939–949.
- Morley JF, Brignull HR, Weyers JJ, Morimoto RI (2002) The threshold for polyglutamine expansion protein aggregation and cellular toxicity is dynamic and influenced by aging in Caenorhabditis elegans. *Proc Natl Acad Sci USA* 99(16):10417–10422.
- Cohen E, Bieschke J, Perciavalle RM, Kelly JW, Dillin A (2006) Opposing activities protect against age-onset proteotoxicity. *Science* 313(5793):1604–1610.
- Lithgow GJ, White TM, Melov S, Johnson TE (1995) Thermotolerance and extended life-span conferred by single-gene mutations and induced by thermal stress. *Proc Natl Acad Sci USA* 92(16):7540–7544.
- Hsu AL, Murphy CT, Kenyon C (2003) Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science* 300(5622):1142–1145.
- Morley JF, Morimoto RI (2004) Regulation of longevity in Caenorhabditis elegans by heat shock factor and molecular chaperones. *Mol Biol Cell* 15(2):657–664.
- Walker GA, Lithgow GJ (2003) Lifespan extension in C. elegans by a molecular chaperone dependent upon insulin-like signals. *Aging Cell* 2(2):131–139.
- Fonte V, et al. (2008) Suppression of in vivo beta-amyloid peptide toxicity by overexpression of the HSP-16.2 small chaperone protein. *J Biol Chem* 283(2):784–791.
- Gengyo-Ando K, Kagawa H (1991) Single charge change on the helical surface of the paramyosin rod dramatically disrupts thick filament assembly in Caenorhabditis elegans. *J Mol Biol* 219(3):429–441.
- MacLeod AR, Waterston RH, Fishpool RM, Brenner S (1977) Identification of the structural gene for a myosin heavy-chain in Caenorhabditis elegans. *J Mol Biol* 114(1):133–140.
- Clark SG, Shurland DL, Meyerowitz EM, Bargmann CI, van der Bliek AM (1997) A dynamin GTPase mutation causes a rapid and reversible temperature-inducible locomotion defect in C. elegans. *Proc Natl Acad Sci USA* 94(19):10438–10443.
- Fares H, Greenwald I (2001) Genetic analysis of endocytosis in Caenorhabditis elegans: Coelomocyte uptake defective mutants. *Genetics* 159(1):133–145.
- Ikeda K, Ozawa S, Hagiwara S (1976) Synaptic transmission reversibly conditioned by single-gene mutation in Drosophila melanogaster. *Nature* 259(5543):489–491.
- Nonet ML (1999) Visualization of synaptic specializations in live C. elegans with synaptic vesicle protein-GFP fusions. *J Neurosci Methods* 89(1):33–40.
- Mawuenyega KG, et al. (2003) Large-scale identification of Caenorhabditis elegans proteins by multidimensional liquid chromatography-tandem mass spectrometry. *J Proteome Res* 2(1):23–35.
- Washburn MP, Wolters D, Yates JR, III (2001) Large-scale analysis of the yeast proteome by multidimensional protein identification technology. *Nat Biotechnol* 19(3):242–247.
- Eisenmann DM, Kim SK (1997) Mechanism of activation of the Caenorhabditis elegans ras homologue let-60 by a novel, temperature-sensitive, gain-of-function mutation. *Genetics* 146(2):553–565.
- Kayser EB, Morgan PG, Hoppel CL, Sedensky MM (2001) Mitochondrial expression and function of GAS-1 in Caenorhabditis elegans. *J Biol Chem* 276(23):20551–20558.
- Hunt-Newbury R, et al. (2007) High-throughput in vivo analysis of gene expression in Caenorhabditis elegans. *PLoS Biol* 5(9):e237.
- Lund J, et al. (2002) Transcriptional profile of aging in C. elegans. *Curr Biol* 12(18):1566–1573.
- McCarroll SA, et al. (2004) Comparing genomic expression patterns across species identifies shared transcriptional profile in aging. *Nat Genet* 36(2):197–204.
- Kim J, Lowe T, Hoppe T (2008) Protein quality control gets muscle into shape. *Trends Cell Biol* 18(6):264–272.
- Kirkwood TB (2008) Understanding ageing from an evolutionary perspective. *J Intern Med* 263(2):117–127.
- Van Voorhies WA, Fuchs J, Thomas S (2005) The longevity of Caenorhabditis elegans in soil. *Biol Lett* 1(2):247–249.
- Dillin A, Crawford DK, Kenyon C (2002) Timing requirements for insulin/IGF-1 signaling in C. elegans. *Science* 298(5594):830–834.
- Rea SL, Wu D, Cypser JR, Vaupel JW, Johnson TE (2005) A stress-sensitive reporter predicts longevity in isogenic populations of Caenorhabditis elegans. *Nat Genet* 37(8):894–898.
- Brunet A, et al. (2004) Stress-dependent regulation of FOXO transcription factors by the SIRT1 deacetylase. *Science* 303(5666):2011–2015.
- Westerheide SD, Anckar J, Stevens SM, Jr, Sistonen L, Morimoto RI (2009) Stress-inducible regulation of heat shock factor 1 by the deacetylase SIRT1. *Science* 323:1063–1066.
- Lee SS, Kennedy S, Tolonen AC, Ruvkun G (2003) DAF-16 target genes that control C. elegans life-span and metabolism. *Science* 300(5619):644–647.
- Wolkow CA, Kimura KD, Lee MS, Ruvkun G (2000) Regulation of C. elegans life-span by insulin-like signaling in the nervous system. *Science* 290(5489):147–150.
- Prahlad V, Cornelius T, Morimoto RI (2008) Regulation of the cellular heat shock response in Caenorhabditis elegans by thermosensory neurons. *Science* 320(5877):811–814.
- Sachidanandam R, et al. (2001) A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* 409(6822):928–933.
- Ng PC, Henikoff S (2006) Predicting the effects of amino acid substitutions on protein function. *Annu Rev Genomics Hum Genet* 7:61–80.
- Cleveland DW, Rothstein JD (2001) From Charcot to Lou Gehrig: Deciphering selective motor neuron death in ALS. *Nat Rev Neurosci* 2(11):806–819.
- Gidalevitz T, Krupinski T, Garcia SM, Morimoto RI (2009) Destabilizing protein polymorphisms in the genetic background direct phenotypic expression of mutant SOD1 toxicity. *PLoS Genet* 5(3):e1000399.
- Belin AC, Westerlund M (2008) Parkinson's disease: A genetic perspective. *FEBS J* 275(7):1377–1383.
- Wexler NS, et al. (2004) Venezuelan kindreds reveal that genetic and environmental factors modulate Huntington's disease age of onset. *Proc Natl Acad Sci USA* 101(10):3498–3503.
- Balch WE, Morimoto RI, Dillin A, Kelly JW (2008) Adapting proteostasis for disease intervention. *Science* 319(5865):916–919.
- Ardizzi JP, Epstein HF (1987) Immunohistochemical localization of myosin heavy chain isoforms and paramyosin in developmentally and structurally diverse muscle cell types of the nematode Caenorhabditis elegans. *J Cell Biol* 105(6 Pt 1):2763–2770.
- Nollen EA, et al. (2004) Genome-wide RNA interference screen identifies previously undescribed regulators of polyglutamine aggregation. *Proc Natl Acad Sci USA* 101(17):6403–6408.
- Henderson ST, Johnson TE (2001) daf-16 integrates developmental and environmental inputs to mediate aging in the nematode Caenorhabditis elegans. *Curr Biol* 11(24):1975–1980.